


Search for an annual modulation
in 3.4 yr of CoGeNT data

J.I. Collar

TAUP 2013, Asilomar


A brief chronology of past CoGeNT results

- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu \beta\beta$ -decay searches).




A brief chronology of past CoGeNT results

- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu \beta\beta$ -decay searches).
- First results (PRL 101 (2008) 251301) in a shallow site eliminated the last region of WIMP parameter space allowed for DAMA/LIBRA within a standard halo model (SHM). This exclusion later confirmed by other searches (e.g., CDMS low-threshold analyses).




A brief chronology of past CoGeNT results

- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu\beta\beta$ -decay searches).
- First results (PRL 101 (2008) 251301) in a shallow site eliminated the last region of WIMP parameter space allowed for DAMA/LIBRA within a standard halo model (SHM). This exclusion later confirmed by other searches (e.g., CDMS low-threshold analyses).
- Irreducible low-energy exponential excess found following surface event rejection (PRL 106 (2011) 131301). WIMP interpretation in vicinity of DAMA/LIBRA ROI. The improved rejection allowed for by larger exposure, and a best-effort at background simulation, have thus far failed to account for this excess (PRD 88 (2013) 012002).




A brief chronology of past CoGeNT results

- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu\beta\beta$ -decay searches).
- First results (PRL 101 (2008) 251301) in a shallow site eliminated the last region of WIMP parameter space allowed for DAMA/LIBRA within a standard halo model (SHM). This exclusion later confirmed by other searches (e.g., CDMS low-threshold analyses).
- Irreducible low-energy exponential excess found following surface event rejection (PRL 106 (2011) 131301). WIMP interpretation in vicinity of DAMA/LIBRA ROI. The improved rejection allowed for by larger exposure, and a best-effort at background simulation, have thus far failed to account for this excess (PRD 88 (2013) 012002). Possible very similar excess in NR band in CDMS-Ge data (Collar & Fields, arXiv:1204.3559, see also R. Nelson's talk later today).



A brief chronology of past CoGeNT results


- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu\beta\beta$ -decay searches).
- First results (PRL 101 (2008) 251301) in a shallow site eliminated the last region of WIMP parameter space allowed for DAMA/LIBRA within a standard halo model (SHM). This exclusion later confirmed by other searches (e.g., CDMS low-threshold analyses).
- Irreducible low-energy exponential excess found following surface event rejection (PRL 106 (2011) 131301). WIMP interpretation in vicinity of DAMA/LIBRA ROI. The improved rejection allowed for by larger exposure, and a best-effort at background simulation, have thus far failed to account for this excess (PRD 88 (2013) 012002). Possible very similar excess in NR band in CDMS-Ge data (Collar & Fields, arXiv:1204.3559, see also R. Nelson's talk later today).
- Run interrupted by Soudan fire: 15 mo of data exhibit low-energy modulation in bulk events, compatible with DAMA/LIBRA (PRL 107 (2011) 141301). Data-sharing allowed for independent analyses and interpretations.



A brief chronology of past CoGeNT results


- CoGeNT employs PPCs (JCAP 09 (2007) 009) to search for low-mass WIMPs, specifically aiming to test the DAMA/LIBRA claim. PPCs offer required stability, low threshold, and rejection of surface events. At higher energies, rejection of gamma backgrounds (MAJORANA and GERDA, $0\nu \beta\beta$ -decay searches).
- First results (PRL 101 (2008) 251301) in a shallow site eliminated the last region of WIMP parameter space allowed for DAMA/LIBRA within a standard halo model (SHM). This exclusion later confirmed by other searches (e.g., CDMS low-threshold analyses).
- Irreducible low-energy exponential excess found following surface event rejection (PRL 106 (2011) 131301). WIMP interpretation in vicinity of DAMA/LIBRA ROI. The improved rejection allowed for by larger exposure, and a best-effort at background simulation, have thus far failed to account for this excess (PRD 88 (2013) 012002). Possible very similar excess in NR band in CDMS-Ge data (Collar & Fields, arXiv:1204.3559, see also R. Nelson's talk later today).
- Run interrupted by Soudan fire: 15 mo of data exhibit low-energy modulation in bulk events, compatible with DAMA/LIBRA (PRL 107 (2011) 141301). Data-sharing allowed for independent analyses and interpretations.
- Much ensuing action: CRESST and CDMS-Si anomalies, XENON exclusions (and criticisms thereof), etc. TBD.

Ge-Si detector landscape (just part of the story)




What is new?

- Detector recovered from 3 mo post-fire outage w/o significant changes in performance. It has been continuously taking data ever since. All data are usable (compare to 10%-40% in CDMS low-energy analyses).





What is new?


- Detector recovered from 3 mo post-fire outage w/o significant changes in performance. It has been continuously taking data ever since. All data are usable (compare to 10%-40% in CDMS low-energy analyses).
- Large exposure allows optimal separation of bulk and surface events down to 0.5 keVee threshold. Rise-time behavior as predicted by simulations and calibrations (PRD 88 (2013) 012002). Smooth variation of fit parameters with energy.



Regions selected for “toy” analysis



M. Bellis
et al.,
in preparation.



See also
poster by
M. Kos.


What is new?

- Detector recovered from 3 mo post-fire outage w/o significant changes in performance. It has been continuously taking data ever since. All data are usable (compare to 10%-40% in CDMS low-energy analyses).
- Large exposure allows optimal separation of bulk and surface events down to 0.5 keVee threshold. Rise-time behavior as predicted by simulations and calibrations (PRD 88 (2013) 012002). Smooth variation of fit parameters with energy.
- Paper under review, preprint to appear soon. Data to be released in energy, time-stamp, and rise-time format. A straightforward analysis indicates a persistent annual modulation exclusively at low energy and for bulk events. Best-fit phase consistent with DAMA/LIBRA (small offset may be meaningful). Similar best-fit parameters to 15 mo dataset, but with much better bulk/surface separation ($\sim 90\%$ SA for $\sim 90\%$ BR)

Dotted: free T
Solid: T = 365 d


See also
poster by M. Kos.

Additional
four months of
unanalyzed
data acquired
(run is still
ongoing)




What is new?

- Detector recovered from 3 mo post-fire outage w/o significant changes in performance. It has been continuously taking data ever since. All data are usable (compare to 10%-40% in CDMS low-energy analyses).
- Large exposure allows optimal separation of bulk and surface events down to 0.5 keVee threshold. Rise-time behavior as predicted by simulations and calibrations (PRD 88 (2013) 012002). Smooth variation of fit parameters with energy.
- Paper under review, preprint to appear soon. Data to be released in energy, time-stamp, and rise-time format. A straightforward analysis indicates a persistent annual modulation exclusively at low energy and for bulk events. Best-fit phase consistent with DAMA/LIBRA (small offset may be meaningful). Similar best-fit parameters to 15 mo dataset, but with much better bulk/surface separation ($\sim 90\%$ SA for $\sim 90\%$ BR)
- Unoptimized frequentist analysis yields $\sim 2.2\sigma$ preference over null hypothesis. This however does not take into account the possible relevance of the modulation amplitude found...




What is new?

- Detector recovered from 3 mo post-fire outage w/o significant changes in performance. It has been continuously taking data ever since. All data are usable (compare to 10%-40% in CDMS low-energy analyses).
- Large exposure allows optimal separation of bulk and surface events down to 0.5 keVee threshold. Rise-time behavior as predicted by simulations and calibrations (PRD 88 (2013) 012002). Smooth variation of fit parameters with energy.
- Paper under review, preprint to appear soon. Data to be released in energy, time-stamp, and rise-time format. A straightforward analysis indicates a persistent annual modulation exclusively at low energy and for bulk events. Best-fit phase consistent with DAMA/LIBRA (small offset may be meaningful). Similar best-fit parameters to 15 mo dataset, but with much better bulk/surface separation ($\sim 90\%$ SA for $\sim 90\%$ BR)
- Unoptimized frequentist analysis yields $\sim 2.2\sigma$ preference over null hypothesis. This however does not take into account the possible relevance of the modulation amplitude found...
- Modulation amplitude is 4-7 times larger than that predicted by the SHM. Finding an absence of modulation would have severely constrained non-standard halo models as explanations for DAMA/LIBRA.



What to make out of this?


- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.



Rough sketch: two WIMPs inducing the same DAMA/LIBRA observable (absolute modulation), but having a different fractional modulation. A SHM cannot induce the large modulation case.

What to make out of this?


- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.



Rough sketch: two WIMPs inducing the same DAMA/LIBRA observable (absolute modulation), but having a different fractional modulation. A SHM cannot induce the large modulation case.

What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.




e.g., [arXiv:1209.3339](https://arxiv.org/abs/1209.3339)

FIG. 3: A comparison of the shapes of the total rate shown at two periods of the year, corresponding to the times of year at which the rate is minimized and maximized, as well as the modulation amplitude, for three different halo components: SHM (left), debris flow (middle), stream (right). The normalization between panels is arbitrary.


What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.
- A large modulation in WIMP rate can arise naturally in many non-SHM, and in particular for large values of v_{\min} , like those probed for $m_\chi \sim 10$ GeV (and small Q_{Na}).



e.g., [arXiv:1209.3339](https://arxiv.org/abs/1209.3339)


FIG. 3: A comparison of the shapes of the total rate shown at two periods of the year, corresponding to the times of year at which the rate is minimized and maximized, as well as the modulation amplitude, for three different halo components: SHM (left), debris flow (middle), stream (right). The normalization between panels is arbitrary.



e.g., [JCAP08\(2012\)027](https://arxiv.org/abs/1209.3339),
[arXiv:1111.0292](https://arxiv.org/abs/1111.0292), [arXiv:1111.0292...](https://arxiv.org/abs/1111.0292)


What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.
- A large modulation in WIMP rate can arise naturally in many non-SHM, and in particular for large values of v_{\min} , like those probed for $m_\chi \sim 10$ GeV (and small Q_{Na}).
- A large fractional modulation for DAMA/LIBRA, corresponding to that found in CoGeNT data, brings it into agreement with other anomalies (CoGeNT, CDMS-Si, and CRESST if slightly underestimating bckgs)




What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.
- A large modulation in WIMP rate can arise naturally in many non-SHM, and in particular for large values of v_{\min} , like those probed for $m_\chi \sim 10$ GeV (and small Q_{Na}).
- A large fractional modulation for DAMA/LIBRA, corresponding to that found in CoGeNT data, brings it into agreement with other anomalies (CoGeNT, CDMS-Si, and CRESST if slightly underestimating bckgs)
- However, this statement ignores the 800 lb gorilla in the room: Q_{Na} is not well-established (whereas CoGeNT's Q_{Ge} is solid, see JCAP 09 (2007) 009).



Full disclosure: I am the author of these two measurements, but was hoping to find $Q_{\text{Na}} \sim 0.4...$ (see arguments in PRD 82 (2010) 123509)




What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.
- A large modulation in WIMP rate can arise naturally in many non-SHM, and in particular for large values of v_{\min} , like those probed for $m_\chi \sim 10$ GeV (and small Q_{Na}).
- A large fractional modulation for DAMA/LIBRA, corresponding to that found in CoGeNT data, brings it into agreement with other anomalies (CoGeNT, CDMS-Si, and CRESST if slightly underestimating bckgs)
- However, this statement ignores the 800 lb gorilla in the room: Q_{Na} is not well-established (whereas CoGeNT's Q_{Ge} is solid, see JCAP 09 (2007) 009).
- Recent efforts to isolate astrophysical (halo) uncertainties (e.g., PRD 83 (2011) 103514) are the best way to examine this complex situation.

Expected modulation amplitude in CoGeNT (upper limit) as a function of Q_{Na} and WIMP mass, taking DAMA/LIBRA as the input, and removing astrophysical uncertainties. Units are the same as in CoGeNT (counts/30d) plot a few transparencies above.

Plot by Chris Kelso, using the halo-independent formalism by P. Fox *et al.* (PRD 83 (2011) 103514, see also PRD 85 (2012) 043515).




What to make out of this?

- Most (uneducated) statements about the incompatibility of DAMA/LIBRA with other current anomalies forget to notice the underlying assumption of a SHM.
- The truth is, DAMA/LIBRA shows no obvious spectral excess, and therefore we cannot know the magnitude of a possible **fractional** modulation in WIMP rate. CoGeNT provides both spectral and modulation information, removing this source of uncertainty.
- Most recent work in halo simulations indicates that finding a SHM (specifically a pure Maxwellian distribution at large v) would be the surprise.
- A large modulation in WIMP rate can arise naturally in many non-SHM, and in particular for large values of v_{\min} , like those probed for $m_\chi \sim 10$ GeV (and small Q_{Na}).
- A large fractional modulation for DAMA/LIBRA, corresponding to that found in CoGeNT data, brings it into agreement with other anomalies (CoGeNT, CDMS-Si, and CRESST if slightly underestimating bckgs)
- However, this statement ignores the 800 lb gorilla in the room: Q_{Na} is not well-established (whereas CoGeNT's Q_{Ge} is solid, see JCAP 09 (2007) 009).
- Recent efforts to isolate astrophysical (halo) uncertainties (e.g., PRD 83 (2011) 103514) are the best way to examine this complex situation.
- The actual few-keVnr value of Q_{Na} will be keystone in determining if DAMA/LIBRA is in agreement with all other low-energy anomalies, or broadly excluded for any WIMP halo model.

Expected modulation amplitude in CoGeNT (upper limit) as a function of Q_{Na} and WIMP mass, taking DAMA/LIBRA as the input, and removing astrophysical uncertainties. Units are the same as in CoGeNT (counts/30d) plot a few transparencies above.

Plot by Chris Kelso, using the halo-independent formalism by P. Fox *et al.* (PRD 83 (2011) 103514, see also PRD 85 (2012) 043515).



If Q_{Na} is the standard ~ 0.3 , then move on, there is nothing to see here...

A few personal impressions:

- If Q_{Na} for 2–6 keVee in NaI[TL] is the usual ~ 0.3 , then DAMA/LIBRA and CoGeNT's observations most probably have nothing to do with each other, not within a WIMP context. It would then seem possible to constraint non-SHM scenarios for DAMA, using CoGeNT data.

A few personal impressions:


- If Q_{Na} for 2–6 keVee in NaI[Tl] is the usual ~ 0.3 , then DAMA/LIBRA and CoGeNT's observations most probably have nothing to do with each other, not within a WIMP context. It would then seem possible to constraint non-SHM scenarios for DAMA, using CoGeNT data.
- If on the other hand this Q_{Na} is ~ 0.15 , then four independent pieces of information may be in agreement: CoGeNT's spectral shape, its modulation, DAMA's modulation, and Q_{Na} (recall, no spectral WIMP info from DAMA). Agreement between all present DM anomalies is an enticing possible outcome.

A few personal impressions:


- If Q_{Na} for 2–6 keVee in NaI[Tl] is the usual ~ 0.3 , then DAMA/LIBRA and CoGeNT's observations most probably have nothing to do with each other, not within a WIMP context. It would then seem possible to constraint non-SHM scenarios for DAMA, using CoGeNT data.
- If on the other hand this Q_{Na} is ~ 0.15 , then four independent pieces of information may be in agreement: CoGeNT's spectral shape, its modulation, DAMA's modulation, and Q_{Na} (recall, no spectral WIMP info from DAMA). Agreement between all present DM anomalies is an enticing possible outcome.
- Clearly, additional measurements of Q_{Na} are in order.

A few personal impressions:

- If Q_{Na} for 2–6 keV in NaI(Tl) is the usual ~ 0.3 , then DAMA/LIBRA and CoGeNT's observations most probably have nothing to do with each other, not within a WIMP context. It would then seem possible to constraint non-SHM scenarios for DAMA, using CoGeNT data.
- If on the other hand this Q_{Na} is ~ 0.15 , then four independent pieces of information may be in agreement: CoGeNT's spectral shape, its modulation, DAMA's modulation, and Q_{Na} (recall, no spectral WIMP info from DAMA). Agreement between all present DM anomalies is an enticing possible outcome.
- Clearly, additional measurements of Q_{Na} are in order.
- It seems feasible to attempt an early exercise in "WIMP astronomy" using existing CoGeNT data, i.e., reverse-engineer the halo properties that would give rise to the observed modulation. We are attempting this, in collaboration with M. Bellis and C. Kelso. These predictions could be tested by GAIA satellite observations very soon.



SLOAN star-count map
showing Milky Way tidal streams




A few personal impressions:

- If Q_{Na} for 2–6 keVee in NaI(Tl) is the usual ~ 0.3 , then DAMA/LIBRA and CoGeNT's observations most probably have nothing to do with each other, not within a WIMP context. It would then seem possible to constraint non-SHM scenarios for DAMA, using CoGeNT data.
- If on the other hand this Q_{Na} is ~ 0.15 , then four independent pieces of information may be in agreement: CoGeNT's spectral shape, its modulation, DAMA's modulation, and Q_{Na} (recall, no spectral WIMP info from DAMA). Agreement between all present DM anomalies is an enticing possible outcome.
- Clearly, additional measurements of Q_{Na} are in order.
- It seems feasible to attempt an early exercise in "WIMP astronomy" using existing CoGeNT data, i.e., reverse-engineer the halo properties that would give rise to the observed modulation. We are attempting this, in collaboration with M. Bellis and C. Kelso. These predictions could be tested by GAIA satellite observations very soon.
- We should not be left forever wondering about XENON-100 excluding this low-mass ROI or not: *in situ* calibrations with the Y/Be source described in PRL 110 (2013) 211101 should settle this issue, once for all. LUX and XMASS results should also cast light (both feature significantly lower thresholds).



PRL 110, 211101 (2013)



Standing challenge to XENON-100:
we hear they will gallantly take it up.

(choose your own exiting quote here)

- “In so far as a scientific statement speaks about reality, it must be falsifiable; and in so far as it is not falsifiable, it does not speak about reality”. K. Popper
- “Everything should be made as simple as possible, but not simpler”. A. Einstein



(choose your own exiting quote here)

- “In so far as a scientific statement speaks about reality, it must be falsifiable; and in so far as it is not falsifiable, it does not speak about reality”. K. Popper
- “Everything should be made as simple as possible, but not simpler”. A. Einstein

(We have not even opened the particle physics can-of-worms today. However, old grandpa Al is very disappointed at you, if you were really expecting the spherical cow)

C-4: coming up very soon

- * First C-4 detector features $\sim 1/3$ of the noise of the existing GoGeNT detector, at $\sim \times 3$ its mass (1.3 kg)
- * Not a one-off: its noise characteristics are now reproducible (CANBERRA R&D supported by NSF award PHY-1003940). Second detector expected to reach the same noise figure at 2 kg, the realistic PPC maximum.
- * C-4 aims at a $\times 10$ total mass increase, $\sim \times 20$ background decrease, and substantial threshold reduction. Soudan is our laboratory of choice, assuming its continuity.




Design and assembly of ULB cryostat at PNNL

Appendix: MALBEK and CoGeNT side-to-side

(apologies, I am a speaker in another session during MALBEK's talk)


P. Finnerty, Ph.D. thesis.



MALBEK is an ideal instrument to test CoGeNT (same detector design and mass, several parts provided by CoGeNT, identical ~160 eV FWHM intrinsic detector noise, similar background achieved)

Appendix: MALBEK and CoGeNT side-to-side

(apologies, I am a speaker in another session during MALBEK's talk)




Unfortunately, poor decisions in the choice of MAJORANA DAQ inject a dominant source of electronic noise. The above shows a comparison for a typical 1.2 keV event in both, pre-denoising, following significant work to reduce polling noise in MALBEK. In a densely-packed MAJORANA-demonstrator array, noise performance can be expected to further degrade.

Appendix: MALBEK and CoGeNT side-to-side


(apologies, I am a speaker in another session during MALBEK's talk)

Adapted from
P. Finnerty, Ph.D. thesis.



99% SA boundaries
for fast pulser signals,
overlapped on
MALBEK's pulser data.

This shift is key.
Dotted line is
my estimated centroid.
(I am hoping this will be
duly emphasized in
MALBEK's talk.
Perhaps I am being naive)




As a result, MALBEK cannot measure preamplifier rise-times below ~ 2 keVee, i.e., at the energies of interest. A variable W_{par} derived from wavelet analysis is seen to correlate with r.t. at high energy, but rapidly becomes the same for surface and bulk events at low energy. MALBEK attempts to exploit W_{par} only down to 0.6 keVee (whereas CoGeNT measures true r.t. down to 0.5 keVee).

Appendix: MALBEK and CoGeNT side-to-side

(apologies, I am a speaker in another session during MALBEK's talk)

Extracted from
P. Finnerty, Ph.D. thesis.



Fair comparison
(subset of CoGeNT
data selected to match
MALBEK's underground
exposure).


Full CoGeNT
dataset (x5 this exposure)
would display an
even crisper surface-bulk
separation
(see upcoming paper).

In the energy ROI ($\sim < 1.5$ keVee) MALBEK's surface and bulk events are essentially indistinguishable, while CoGeNT preserves the ability to separate these down to threshold. Keeping in mind that the modulation observed in CoGeNT is a $\sim 5\%$ oscillation of the overall (surface + bulk) rate, a search for an annual modulation in MALBEK seems a futile exercise.


As a former MAJORANA collaborator, I should express my concern about the impact of DAQ hardware choices on its low-energy physics potential. See PRL 101 (2008) 251301 for a discussion of MAJORANA's potential as a low-mass WIMP detector.

Appendix: MALBEK and CoGeNT side-to-side

(apologies, I am a speaker in another session during MALBEK's talk)



Trying to squeeze blood out of a turnip: an aggressive W_{par} cut runs the risk of throwing out baby and bathwater at low-E.



The larger statistics of CoGeNT (x5 MALBEK's) and much better surface-bulk separation produce little dependence of spectral shape and limits on choice of rise-time cuts. Quite the opposite for MALBEK (order of magnitude difference).


The low-E W_{par} distribution in MALBEK (see above) strongly points at roughly the same bulk and surface event contributions. In other words, a markedly exaggerated exclusion limit when applying aggressive W_{par} cuts.

MALBEK would do well in waiting to understand their low-E separation between surface and bulk events. This has been very beneficial for CoGeNT.

Reserve

Fair comparison

(same grayscale and number of events plotted)



CDMS-II data are now publicly available on ArXiv.